

Date: June 29, 2005

Contacts: Vanee Vines, Senior Media Relations Officer

Megan Petty, Media Relations Assistant

Office of News and Public Information

202-334-2138; e-mail <news@nas.edu>

FOR IMMEDIATE RELEASE

Low Levels of Ionizing Radiation May Cause Harm

WASHINGTON -- A preponderance of scientific evidence shows that even low doses of ionizing radiation, such as gamma rays and X-rays, are likely to pose some risk of adverse health effects, says a new report from the

radiation can be demonstrated to be harmless or beneficial," said committee chair Richard R. Monson, associate dean for professional education and professor of epidemiology, Harvard School of Public Health, Boston. "The health risks – particularly the development of solid cancers in organs – rise proportionally with exposure. At low doses of radiation, the risk of inducing solid cancers is very small. As the overall lifetime exposure increases, so does the risk." The report is the seventh in a series on the biological effects of ionizing radiation.

Assessing Health Risks

The committee's risk models for exposure to low-level ionizing radiation were based on a sex and age distribution similar to that of the entire U.S. population, and refer to the risk that an individual would face over his or her life span. These models predict that about one out of 100 people would likely develop solid cancer or leukemia from an exposure of 0.1 Sv (100 mSv). About 42 additional people in the same group would be expected to develop solid cancer or leukemia from other causes. Roughly half of these cancers would result in death. These particular estimates are uncertain, however, because of limitations in the data used to develop risk models.

Survivors of atomic bombings in Hiroshima and Nagasaki, Japan, were the primary sources of data for estimating risks of most solid cancers and leukemia from exposure to ionizing radiation. The committee's review included an examination of updated cancer-incidence data from tumor registries of the survivors, and of research data on solid cancer deaths -- which is now more abundant because the number of deaths available for analysis has nearly doubled since the Research Council published its previous report on this topic in 1990. The committee combined this information with data on people who had been medically exposed to radiation to estimate risks of breast cancer in women and thyroid cancer. Data from additional medical studies and from studies of people exposed to radiation through their occupations also were evaluated and found to be compatible with the committee's statistical models. Follow-up studies should continue for the indefinite future, the report says.

Adverse hereditary health effects that could be attributed to radiation have not been found in studies of children whose parents were exposed to radiation from the atomic bombs. However, studies of mice and other organisms have produced extensive data showing that radiation-induced cell mutations in sperm and eggs can be passed on to offspring, the report says. There is no reason to believe that such mutations could not also be passed on to human offspring. The failure to observe such effects in Hiroshima and Nagasaki probably reflects an insufficiently large survivor population.

Follow-up studies of people who receive computed tomography (CT) scans, especially children, should be conducted, the report adds. Also needed are studies of infants who are exposed to diagnostic radiation because catheters have been placed in their hearts, as well as infants who receive multiple X-rays to monitor pulmonary development. CT scans, often referred to as whole body scans, result in higher doses of radiation than typically experienced with conventional X-rays.

Sources of Ionizing Radiation

People are exposed to natural background ionizing radiation from the universe, the ground, and basic activities such as eating, drinking, and breathing. These sources account for about 82 percent of human exposure.

Nationwide, man-made radiation comprises 18 percent of human exposure. In this overall category, medical X-rays and nuclear medicine account for about 79 percent, the report says. Elements in consumer products -- such as tobacco, tap water, and building materials -- account for another 16 percent. Occupational exposure, fallout, and the use of nuclear fuel constitute roughly 5 percent of the man-made component nationwide.

Factors that could increase exposure include greater use of radiation for medical purposes, working around radioactive materials, and smoking tobacco. Living at low altitudes, where there is less cosmic radiation, and living and working on the upper floors of buildings, where there is less radon gas -- a primary source of natural ionizing radiation -- are factors that could decrease exposure.

The report was sponsored by the U.S. departments of Defense, Energy, and Homeland Security, the U.S. Nuclear Regulatory Commission, and the U.S. Environmental Protection Agency. The National Research Council is the principal operating arm of the National Academy of Sciences and the National Academy of Engineering. It is a private, nonprofit institution that provides science and technology advice under a congressional charter. A committee roster follows.

Copies of [Health Risks from Exposure to Low Levels of Ionizing Radiation \(BEIR VII - Phase 2\)](#) will be available this summer from the National Academies Press; tel. 202-334-3313 or 1-800-624-6242 or on the Internet at <http://www.nap.edu>. Reporters may obtain a copy from the Office of News and Public Information (contacts listed above).

[This news release and report are available at <http://national-academies.org>]

NATIONAL RESEARCH COUNCIL

Division on Earth and Life Studies

Board on Radiation Effects Research

Committee to Assess Health Risks From Exposure to Low Levels of Ionizing Radiation

Richard R. Monson, M.D., Sc.D. (chair)

Associate Dean for Professional Education, and
Professor of Epidemiology
School of Public Health
Harvard University
Boston

James E. Cleaver, Ph.D.(1) (vice chair)
Professor of Dermatology
Cancer Center and Department of Pharmaceutical Chemistry
University of California
San Francisco

Herbert L. Abrams, M.D. (2)
Professor Emeritus of Radiology
Stanford University Medical School, and
Member in Residence
Stanford Center for International Security and Cooperation
Stanford, Calif.

Eula Bingham, Ph.D. (2)
Professor of Environmental Health
University of Cincinnati
Cincinnati

Patricia A. Buffler, Ph.D. (2)
Kenneth and Marjorie Kaiser Chair of Cancer Epidemiology, and
Professor of Epidemiology
School of Public Health
University of California
Berkeley

Elisabeth Cardis, Ph.D.
Chief, Unit of Radiation and Cancer
International Agency for Research on Cancer
Lyon, France

Roger Cox, Ph.D.
Director
National Radiological Protection Board

Chilton, United Kingdom

Scott Davis, Ph.D.

Professor and Chair

Department of Epidemiology

School of Public Health and Community Medicine

University of Washington, and

Full Member

Program in Epidemiology

Division of Public Health Sciences

Fred Hutchinson Cancer Research Center

Seattle

William C. Dewey, Ph.D.

Emeritus Professor of Radiation Oncology

University of California

San Francisco

Ethel S. Gilbert, Ph.D.

Biostatistician

Radiation Epidemiology Branch

National Cancer Institute

Bethesda, Md.

Albrecht Kellerer, Ph.D.

Professor Emeritus

University of Munich

Munich, Germany

Daniel Krewski, Ph.D., M.H.A.

Director

McLaughlin Centre for Population Health Risk Assessment, and

Professor of Medicine and of Epidemiology and Community Medicine

University of Ottawa

Ontario, Canada

Tomas Lindahl, M.D.

Director

Clare Hall Laboratories

Cancer Research U.K.
London

Katherine E. Rowan, Ph.D.

Professor and Associate Chair
Department of Communication
George Mason University
Fairfax, Va.

K. Sankaranarayanan, Ph.D.

Professor Emeritus
Department of Toxicogenetics
Leiden University Medical Centre
Leiden, Netherlands

Daniel W. Schafer, Ph.D.

Professor
Department of Statistics
Oregon State University
Corvallis

Robert L. Ullrich, Ph.D.

Barbara Cox Anthony University Chair in Oncology
Departments of Environmental and Radiological Health Sciences and of Clinical Sciences
Colorado State University
Fort Collins

RESEARCH COUNCIL STAFF

Rick Jostes, Ph.D.

Study Director

Evan B. Double, Ph.D.

Director
Board on Radiation Effects Research

(1) Member, National Academy of Sciences

(2) Member, Institute of Medicine